



Cambridge International AS & A Level

CANDIDATE NAME							
CENTRE NUMBER				CANDIDAT NUMBER	E		

BIOLOGY 9700/51

Paper 5 Planning, Analysis and Evaluation

May/June 2025

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.


INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 30.
- The number of marks for each question or part question is shown in brackets [].

This document has 16 pages. Any blank pages are indicated.





1 Pepsin is an enzyme that is present in gastric juice. Gastric juice is secreted into the stomach of humans and many other animals.

2

Pepsin catalyses the hydrolysis of proteins into peptides, as shown in Fig. 1.1.




Fig. 1.1

Egg albumen (egg white) contains a high proportion of protein.

10% albumen solution is a cloudy-white colour. This becomes colourless when pepsin is added to the 10% albumen solution.

A student used a colorimeter to follow the progress of the hydrolysis of protein by pepsin.

(a) (i) The student used a blue filter in the colorimeter.

Outline one other step the student should carry out to prepare the colorimeter so the correct measurements of absorbance can be obtained.	ıat
	[1]

- (ii) After preparing the colorimeter, the student:
 - made a pH2.0 albumen solution by mixing 10 cm³ of 10% albumen solution with 10 cm³ of pH2.0 buffer solution
 - added 3.0 cm³ of the pH2.0 albumen solution to a colorimeter tube
 - added a small volume of pepsin solution to the colorimeter tube
 - immediately placed the colorimeter tube in the colorimeter
 - recorded the absorbance of the mixture every 20 seconds for 5 minutes
 - calculated the rate of change in absorbance as a measure of the rate of protein hydrolysis.

The results are shown in Fig. 1.2.



/au






Fig. 1.2

Use Fig. 1.2 to calculate the rate of change in absorbance at pH2.0 between 40s and 120s.

Show your working and include appropriate units.

(b) The student decided to investigate the effect of pH on the rate of protein hydrolysis by pepsin.

From the internet the student found out that pepsin is inactive at pH6.5 and above.

(i) The student was provided with a colorimeter and standard laboratory apparatus.

Describe a method that the student could use to investigate the effect of pH on the rate of protein hydrolysis by pepsin **and** to determine the optimum pH of pepsin.

Your method should be set out in a logical order and be detailed enough to allow another person to follow it.

Details of now to prepare and use the colonimeter should not be included.
[7]



(ii) Complete the sketch graph in Fig. 1.3 to predict the effect of pH on the rate of protein hydrolysis by pepsin.

Include a label for the y-axis in your answer.




Fig. 1.3

[2]



c) (i) Laryngopharyngeal reflux (LPR) is a disease that occurs when gastric juice from the stomach moves up the oesophagus and into the larynx, as shown in Fig. 1.4.




Fig. 1.4

Gastric juice can damage the laryngeal epithelium. Gastric juice contains hydrochloric acid and pepsin.

Some scientists identified a reduction in the quantity of two proteins, CA3 and Sep70, present in the laryngeal epithelium of people with LPR.

The scientists obtained 6 sections of mammalian laryngeal epithelium to investigate how different test conditions affect the quantity of CA3 and Sep70 in the laryngeal epithelium.

Each section of laryngeal epithelium was exposed to different test conditions for 20 minutes, as shown in Table 1.1.

Pepsin is inactive in the presence of the inhibitor, pepstatin.



Table 1.1

treatment used	test condition							
treatment used	1	2	3	4	5	6		
pepsin	yes	yes	yes	yes	no	no		
pepstatin (inhibitor)	no	no	yes	yes	yes	yes		
pH buffer	7.4	4.0	7.4	4.0	7.4	4.0		

Proteins were then extracted from the sections of laryngeal epithelium. The proteins were separated using electrophoresis. CA3 and Sep70 were identified using fluorescent antibodies.

Fig. 1.5 shows the results of the protein electrophoresis.





Fig. 1.5



[Total: 17]

i) The results of this investigation were published in a scientific paper.

8

A studen	t who	read	the	scientific	paper	concluded	that	pepsin	causes	damage	to	the
laryngeal	epith	elium	in pe	eople with	LPR.							

Suggest why the results of this investigation might not support this conclusion.	
	2

* 000080000009 *

9

2 Golden orb weaver spiders, *Nephila pilipes*, are found in East Asia, South-east Asia and Australia. They are active in the day and at night. Golden orb weaver spiders build webs in trees and shrubs to catch different species of insects for food.

All female golden orb weaver spiders are black with yellow spots on their legs and body. Biologists think that this adaptation evolved due to natural selection. Spiders with yellow spots may be able to attract more insects to their webs.

Fig. 2.1 shows a female golden orb weaver spider sitting on a web.



Fig. 2.1

Some biologists visited a forest in East Asia in July 2008. The biologists found five webs that were approximately the same size. Each web belonged to a female golden orb weaver spider.

The biologists decided to investigate how the colour and pattern of spots on the spiders affect the number of insects attracted per hour to each web (insect attraction rate).

The biologists made two-dimensional (2D) models of spiders, as shown in Table 2.1.

10

Table 2.1

model	description	diagram
A	 black body with yellow spots black legs with yellow spots same colour and pattern of spots as golden orb weaver spiders 	
В	 black body with blue spots black legs with blue spots same pattern of spots as golden orb weaver spiders 	
С	 black body with large yellow spot black legs total area of yellow colour is the same as model A 	
D	yellow bodyyellow legs	
E	black body black legs	



At the first web, the biologists:

- removed the living golden orb weaver spider from the web
- placed model A in the centre of the web
- placed a video camera 1 m from the web
- filmed the web and model for 6 hours
- recorded an insect attraction event whenever an insect flew towards the model, touched the model, or touched the web

11

calculated the insect attraction rate of model A.

The procedure was repeated by placing models **B**, **C**, **D** and **E** on the four other webs.

The whole investigation was repeated 25 times, and a mean insect attraction rate for each model was calculated.

(a)	(i)	Identify the dependent variable in this investigation.
		[1]
	(ii)	The pattern of spots on models ${\bf A}$ and ${\bf B}$ was the same as the pattern of spots on female golden orb weaver spiders.
		Identify two other variables that the biologists should standardise when making the models for this investigation.
		[2]
	(iii)	Suggest a suitable control for this investigation.
		[1]





(b) Fig. 2.2 shows the mean insect attraction rates for the five models. The error bars show ± one standard error (SE).




Fig. 2.2

Use the information shown in Table 2.1 and Fig. 2.2 to discuss the conclusions that can be made about the effect of colour and pattern of spots on the models on insect attraction rates.

(c)



(ii) The biologists decided to analyse the data in Fig. 2.2 using *t*-tests.

13

State a null hypothesis the biologists could make before carrying out the <i>t</i> -tests. [1] (iii) State two reasons why the <i>t</i> -test is suitable for analysing the data shown in Fig. 2.2.
[1]
[1]
(iii) State two reasons why the <i>t</i> -test is suitable for analysing the data shown in Fig. 2.2.
(, class the reaction may are a second canadate for amany and add and another many age and
[2]
A student thought that the investigation did not provide enough information about how the colour and pattern of spots on female golden orb weaver spiders help to attract insects to their webs.
Suggest how this investigation could be improved to increase confidence in the results.
[2]

BLANK PAGE



BLANK PAGE



BLANK PAGE

The boundaries and names shown, the designations used and the presentation of material on any maps contained in this question paper/insert do not imply official endorsement or acceptance by Cambridge Assessment International Education concerning the legal status of any country, territory, or area or any of its authorities, or of the delimitation of its frontiers or boundaries.

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

