



Cambridge O Level

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

1171600756

CHEMISTRY 5070/22

Paper 2 Theory May/June 2025

1 hour 45 minutes

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.

This document has 16 pages.

[Total: 5]



1 Choose from the following salts to answer the questions.

aluminium sulfate

2

barium chloride

copper(II) nitrate

copper(II) sulfate

magnesium chloride

potassium iodide

potassium manganate(VII)

silver chloride

sodium bromide

Each salt can be used once, more than once or not at all.

State which salt:

(a)	is prepared using a precipitation reaction	
		[1]
(b)	gives a yellow flame test colour	
		[1]
(c)	dissolves to form a dark purple aqueous solution	
		[1]
(d)	has an aqueous solution that reacts with dilute sulfuric acid to give a white precipitate	
		[1]
(e)	has an aqueous solution that reacts with aqueous bromine.	
		[1]

- 2 A dilute aqueous solution of magnesium chloride is electrolysed using graphite electrodes.
 - (a) Graphite has a high melting point and is inert.

	(i)	Explain why graphite has a high melting point.	
		Use ideas about structure and bonding.	
			. [2]
	(ii)	State one other property of graphite that makes it suitable for use as an electrode du electrolysis.	ıring
			. [1]
(b)		dict the products of the electrolysis of dilute aqueous magnesium chloride with grap ctrodes.	ohite
	prod	duct at anode	
	prod	duct at cathode	[2]
(c)		ten aluminium oxide is electrolysed using graphite electrodes to form oxygen minium.	
	Con	nstruct the ionic half-equation for the reaction at each electrode.	
	read	ction at anode	
	read	ction at cathode	
			[2]
(d)	A m	netal object is electroplated with copper.	
	The	e metal object is the cathode during this electrolysis.	
	Stat	te the name of the substance used for the anode and for the electrolyte.	
	ano	de	
	elec	ctrolyte	[2]
			[4.

[Total: 9]

The equation for the reaction between ethene and bromine is shown in Fig. 3.1.




Fig. 3.1

(a)	Describe the observation when etherie gas is pubbled through addedus brothine.

(b) Table 3.1 shows some bond energies.

Table 3.1

bond	bond energy in kJ/mol	
С—Н	410	
с—с	350	
C=C	610	
Br—Br	193	
C—Br	280	

Show by calculation that the enthalpy change of the reaction between ethene and bromine, ΔH , is $-107\,\mathrm{kJ/mol}$.

[3]

(c) Complete the reaction pathway diagram in Fig. 3.2 for the reaction between ethene and bromine.

Label the:

- reactants
- product
- enthalpy change of the reaction, ΔH
- activation energy, E_a.




Fig. 3.2

how the electronic configuration in a malegula of others

(d) Draw a dot-and-cross diagram to show the electronic configuration in a molecule of ethene.
Show only the outer shell electrons.

[2]

[3]

[Total: 9]



4 Ethanoic acid, CH₃COOH, is a member of the homologous series of carboxylic acids.

Give your answer to **two** significant figures.

(a)	Give	e the general formula of the homologous series of carboxylic acids.	
			[1]
(b)		e characteristic of a homologous series is that all the compounds share similar chemic perties.	cal
	Ехр	lain why the compounds share similar chemical properties.	
			[1]
(c)	The	equation for the reaction between calcium and dilute ethanoic acid is shown.	
		$Ca + 2CH_{3}COOH \to H_{2} + Ca(CH_{3}COO)_{2}$	
	(i)	State the name of the compound Ca(CH ₃ COO) ₂ .	
			[1]
	(ii)	A sample of 1.35 g of calcium is added to excess dilute ethanoic acid.	

Calculate the volume of hydrogen formed measured at room temperature and pressure.

	volume = dm ³ [3]
(d)	Dilute ethanoic acid is a component of vinegar.
	Describe the manufacture of vinegar.
	Include the reactants and conditions.
	rei

(i)

(ii)

7

(e) Butanoic acid and propanoic acid are two other carboxylic acids.

State the molecular formula of butanoic acid.	
	[1]
Draw the displayed formula of propanoic acid.	

		[1]
(iii)	Solid sodium carbonate is added to dilute propanoic acid.	
	Predict an observation for this reaction.	
		[1]
(iv)	Aqueous sodium hydroxide is added to dilute butanoic acid.	
	State the names of the products of this reaction.	
		[1]

[Total: 13]



- 5 The combustion of fossil fuels is used in some power stations.
 - (a) Some power stations use diesel oil as a fuel.

((1)	One compound in diesel oil has the formula $C_{12}H_{26}$.
		Construct the symbol equation to show the complete combustion of $\mathrm{C}_{12}\mathrm{H}_{26}$.
,,		[1]
(1	ii)	The complete combustion of C ₁₂ H ₂₆ produces an air pollutant.
		State one adverse effect of this pollutant.
(ii	ii)	Explain how diesel oil is separated from petroleum.
		[3]
		ur dioxide is removed from the emissions from a power station using calcium carbonate der.
((i)	State one adverse effect of sulfur dioxide as an air pollutant.
		[1]
(i	ii)	One mole of sulfur dioxide reacts with one mole of calcium carbonate to make one mole of carbon dioxide and only one other product.
		Suggest the formula of this product.
		formula [1]

(iii)

State and explain the effect of increasing the temperature on the rate of this reaction.



(iv) Lumps of calcium carbonate are used instead of calcium carbonate powder.

9

State and explain the effect of this change on the rate of this reaction.
[2]

[Total: 11]



- Bromine, Br₂, is in Group VII of the Periodic Table.
 - (a) The melting point of bromine is -7 °C and the boiling point is 59 °C.

(i)	Explain why	bromine is a	a solid at –50°C.	

 [1]

(ii)

Describe the arrangement and motion of bromine molecules at -50 °C.
[3]

(b) A sample of bromine liquid contains 2.408×10^{25} molecules.

One mole of bromine liquid contains 6.02×10^{23} molecules.

Calculate the mass of this sample of bromine liquid.

(c) The ionic equation shows the reaction of bromine with warm concentrated aqueous sodium hydroxide.

$$3Br_2 + 6OH^- \rightarrow BrO_3^- + 5Br^- + 3H_2O$$

State the oxidation number of bromine in Br₂ and in Br⁻.

[2]

(ii) During the reaction bromine is reduced.

Explain why, using ideas about electrons.

* 0000800000011 *

(d) Bromine reacts with hydrogen in a closed system to form an equilibrium mixture. The forward reaction releases thermal energy into the surroundings.

$$Br_2(g) + H_2(g) \rightleftharpoons 2HBr(g)$$

11

	$Br_2(g) + H_2(g) \leftarrow 2HBr(g)$
(i)	The temperature of the equilibrium mixture is increased. The pressure remains constant
	State and explain what happens to the position of equilibrium.
	statement
	explanation
	[2
(ii)	The pressure of the equilibrium mixture is increased. The temperature remains constant
	State and explain what happens to the position of equilibrium.
	statement
	explanation
	·

(e) A bromide of phosphorus contains 7.2% phosphorus by mass.

Calculate the empirical formula of this bromide.

Show your working.

empirical formula[3]

[Total: 16]



7 Iron is used to make stainless steel.

Stai	nless steel is used to make cutlery because it is resistant to rusting.
(a)	Give one other reason why stainless steel is used to make cutlery.
	[1]
(b)	Stainless steel is an alloy.
	Give the meaning of the term alloy.
	[1]
(c)	Iron nails are galvanised with zinc to prevent rusting.
	Explain two ways in which galvanising iron nails with zinc prevents rusting.
	[3]
(d)	Iron reacts with hot dilute sulfuric acid to form hydrogen and aqueous iron(III) sulfate as the only products.
	Construct a symbol equation for this reaction.
	Include state symbols.
	[2]
	[Total: 7]

12

* 0000800000013 *

Fig. 8.1 shows the displayed formula of compound A.

13

Fig. 8.1

(i)	Explain why compound A is saturated.	
		[1]
(ii)	Explain why compound A is an alcohol.	
		[1]
(iii)	Explain why compound A is an ester.	
		[1]

(b) Fig. 8.2 shows compound B.

Fig. 8.2

Explain why compound **A** and compound **B** are a pair of structural isomers.

(c) The displayed formula of compound C is shown in Fig. 8.3.

Fig. 8.3

Compound C is an ester.

Compound **C** is made by the reaction of an alcohol and a carboxylic acid in the presence of a catalyst.

(i)	State the	name of the	type of	catalyst	used in	this	reaction
-----	-----------	-------------	---------	----------	---------	------	----------

- 47
111


(ii) State the name and draw the displayed formula of the alcohol used in this reaction.

displayed formula

[2]

15

(d) Fig. 8.4 shows two monomers that react to make a condensation polymer.



monomer **D**

monomer E

Fig. 8.4

(i) Draw the structure of **one** repeat unit of this condensation polymer.

[2]

(ii) An equal number of moles of the monomers D and E are reacted to make the condensation polymer.

The total mass of the monomers **D** and **E** is 80 g. There is a 100% yield.

Explain why the mass of the condensation polymer made is less than 80 g.

 	 	 	 	[1]

[Total: 10]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.





The Periodic Table of Elements

	III/	2	He	helium 4	10	Ne	neon 20	18	Ā	argon 40	36	궃	krypton 84	54	Xe	xenon 131	98	R	radon	118	Og	oganesson	ı
	\				6	Щ	fluorine 19	17	Cl	chlorine 35.5	35	Ŗ	bromine 80	23	П	iodine 127	82	Αŧ	astatine -	117	<u>S</u>	tennessine	ı
	>				80	0	oxygen 16	16	ഗ	sulfur 32	34	Se	selenium 79	52	<u>e</u>	tellurium 128	84	Ъ	molodium –	116	۲	livermorium	ı
	>				7	z	nitrogen 14	15	凸	phosphorus 31	33	As	arsenic 75	51	Sp	antimony 122	83	Ξ	bismuth 209	115	Mc	moscovium	ı
	≥				9	O	carbon 12	14	S	silicon 28	32	Ge	germanium 73	20	Sn	tin 119	82	Ър	lead 207	114	ŀΙ	flerovium	I
	=				2	Ф	boron 11	13	Αl	aluminium 27	31	Ga	gallium	49	In	indium 115	81	1L	thallium 204	113	R	nihonium	ı
											30	Zu	zinc	48	ρO	cadmium 112	80	Hg	mercury 201	112	C	copemicium	ı
											29	Cn	copper 64	47	Ag	silver 108	62	Au	gold 197	111	Rg	roentgenium	ı
Group											28	Z	nickel	46	Pd	palladium 106	78	പ	platinum 195	110	Ds	darmstadtium	ı
Gre											27	ဝိ	cobalt 59	45	몬	rhodium 103	77	'n	iridium 192	109	Ĭ	meitnerium	ı
		-	エ	hydrogen 1							26	Fe	iron	44	Ru	ruthenium 101	92	SO	osmium 190	108	Hs	hassium	ı
											25	Mn	manganese 55	43	ည	technetium -	75	Re	rhenium 186	107	Bh	bohrium	ı
					_	loq	ass				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	≥	tungsten 184	106	Sg	seaborgium	ı
				Key	atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	5 4	Q N	niobium 93	73	Та	tantalum 181	105	ОР	dubnium	ı
						atc	re				22	F	titanium 48	40	Zr	zirconium 91	72	Ξ	hafnium 178	104	峜	rutherfordium	ı
											21	Sc	scandium 45	39	>	yttrium 89	57–71	lanthanoids		89–103	actinoids		
	=				4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	88	S	strontium 88	56	Ba	barium 137	88	Ra	radium	1
	_				က	:=	lithium 7	=	Na	sodium 23	19	¥	potassium	37	8	rubidium 85	55	S	caesium 133	87	Ŧ	francium	1
					_			_			_						_					_	_

16

71	ר	lutetium	175	103	ئ	lawrencium	ı
70	ΥÞ	ytterbium	173	102	8	nobelium	I
69	H	thulium	169	101	Md	mendelevium	ı
89	ш	erbium	167	100	FB	ferminm	ı
29	웃	holmium	165	66	Es	einsteinium	ı
99	ò	dysprosium	163	86	ర	californium	ı
99	Д	terbium	159	97	Ř	berkelium	ı
64	Вd	gadolinium	157	96	Cm	curium	ı
63	Ш	europium	152	92	Am	americium	I
62	Sm	samarium	150	94	Pu	plutonium	ı
61	Pm	promethium	ı	93	ď	neptunium	ı
09	PΝ	neodymium	144	92	\supset	uranium	238
69	Ā	praseodymium	141	91	Ра	protactinium	231
58	Ce	cerium	140	06	Т	thorium	232
22	Га	lanthanum	139	88	Ac	actinium	ı

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).

