

Cambridge IGCSE™

ADDITIONAL MATHEMATICS**0606/11**

Paper 1

October/November 2025

MARK SCHEME

Maximum Mark: 80

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2025 series for most Cambridge IGCSE, Cambridge International A and AS Level components, and some Cambridge O Level components.

This document consists of **12** printed pages.

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptions for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Mathematics-Specific Marking Principles

- 1 Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
- 2 Unless specified in the question, non-integer answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
- 3 Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
- 4 Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
- 5 Where a candidate has misread a number or sign in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 A or B mark for the misread.
- 6 Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

Annotations guidance for centres

Examiners use a system of annotations as a shorthand for communicating their marking decisions to one another. Examiners are trained during the standardisation process on how and when to use annotations. The purpose of annotations is to inform the standardisation and monitoring processes and guide the supervising examiners when they are checking the work of examiners within their team. The meaning of annotations and how they are used is specific to each component and is understood by all examiners who mark the component.

We publish annotations in our mark schemes to help centres understand the annotations they may see on copies of scripts. Note that there may not be a direct correlation between the number of annotations on a script and the mark awarded. Similarly, the use of an annotation may not be an indication of the quality of the response.

The annotations listed below were available to examiners marking this component in this series.

Annotations

Annotation	Meaning
A	More information required
A0	Accuracy mark awarded zero
A1	Accuracy mark awarded one
A2	Accuracy mark awarded two
A3	Accuracy mark awarded three
B0	Independent mark awarded zero
B1	Independent mark awarded one
B2	Independent mark awarded two
B3	Independent mark awarded three
BOD	Benefit of the doubt
C	Communication mark
X	Incorrect
FT	Follow through
Highlighter	Highlight a key point in the working
ISW	Ignore subsequent work
M0	Method mark awarded zero
M1	Method mark awarded one
M2	Method mark awarded two

Annotation	Meaning
M3	Method mark awarded three
MR	Misread
O	Omission
Off-page comment	Allows comments to be entered at the bottom of the RM marking window and then displayed when the associated question item is navigated to.
On-page comment	Allows comments to be entered in speech bubbles on the candidate response.
Pre	Premature rounding/approximation
SC	Special case
SEEN	Indicates that work/page has been seen
TE	Transcription error
	Correct
XP	Correct answer from incorrect working

MARK SCHEME NOTES

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

M Method marks, awarded for a valid method applied to the problem.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. For accuracy marks to be given, the associated Method mark must be earned or implied.

B Mark for a correct result or statement independent of Method marks.

When a part of a question has two or more ‘method’ steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. The notation ‘dep’ is used to indicate that a particular M or B mark is dependent on an earlier mark in the scheme.

Abbreviations

awrt	answers which round to
cao	correct answer only
dep	dependent
FT	follow through after error
isw	ignore subsequent working
nfww	not from wrong working
oe	or equivalent
rot	rounded or truncated
SC	Special Case
soi	seen or implied

Question	Answer	Marks	Guidance
1(a)	3	B1	
	180° or π	B1	Allow without degrees sign
1(b)	Correct curve 	3	B1 for amplitude 3 B1 for period 180° B1 for correct end points

Question	Answer	Marks	Guidance
2	$x^2 + 2x = kx - 3$ $[\Rightarrow x^2 + (2-k)x + 3 = 0]$	M1	
	$(2-k)^2 - 12 \neq 0$	M1	Dep on previous M mark *Allow use of any inequality or equality
	$k^2 - 4k - 8 \neq 0$	A1	
	$k = \frac{4 \pm \sqrt{4^2 + 4 \times 8}}{2}$ oe	M1	Solve their 3-term quadratic equation in k
	$k = 2 \pm \sqrt{12}$ or $\frac{4 \pm \sqrt{48}}{2}$ oe	A1	
	$k < 2 - 2\sqrt{3}$, $k > 2 + 2\sqrt{3}$ oe	A1	Allow $\sqrt{12}$ for $2\sqrt{3}$
3(a)	$\log_2 x - 4 = 5 \frac{\log_2 2}{\log_2 x}$ or $= \frac{1}{\log_2 x}$	M1	Correct use of change of base
	$(\log_2 x)^2 - 4\log_2 x - 5 = 0$	A1	Arrange as 3-term quadratic equation in $\log_2 x$
	$(\log_2 x - 5)(\log_2 x + 1) = 0$ or $\log_2 x = \frac{4 \pm \sqrt{4^2 - 4 \times (-5)}}{2}$	M1	Dep on previous M mark
	$\log_2 x = 5$ or -1	A1	For either
	$x = 32$ or $x = \frac{1}{2}$	A1	Both
	Alternative method		
	$\frac{\log_x x}{\log_x 2} - 4 = 5 \log_x 2$	(M1)	
	$(5 \log_x 2)^2 + 4 \log_x 2 - 1 = 0$	(A1)	
	$(5 \log_x 2 - 1)(\log_x 2 + 1) = 0$ or $\log_x 2 = \frac{-4 \pm \sqrt{4^2 - 4 \times (-5)}}{10}$	(M1)	Dep on previous M mark
	$\log_x 2 = \frac{1}{5}$ or -1	(A1)	For either
	$x = 32$ or $x = \frac{1}{2}$	(A1)	Both

Question	Answer	Marks	Guidance												
3(b)	$e^{x^2-3-(7-x^2)} = 25$ oe	M1	Arrange as equation with one power of e												
	$2x^2 - 10 = \ln 25$	A1													
	$x^2 = \frac{\ln 25 + 10}{2}$	A1													
	$x = \pm \sqrt{\frac{\ln 25 + 10}{2}}$ oe or $\pm \sqrt{\ln 5 + 5}$	A1	isw												
4(a)	Many-many (mapping) oe or A vertical line meets graph more than once (or twice) oe	B1													
4(b)	<table border="1" style="display: inline-table; vertical-align: middle;"> <tr><td></td><td>✓</td><td></td></tr> <tr><td>✓</td><td></td><td>✓</td></tr> <tr><td></td><td>✓</td><td></td></tr> <tr><td>✓</td><td></td><td></td></tr> </table>		✓		✓		✓		✓		✓			4	B1 for each correct row
	✓														
✓		✓													
	✓														
✓															
4(c)(i)	$\sin x \geq \frac{1}{2}$	M1	May be implied. Allow >												
	$a = 150$ seen	A1													
4(c)(ii)	$0 \leq gf(x) \leq \frac{1}{\sqrt{2}}$	2	B1 for each correct value with correct sign associated with it												
4(c)(iii)	No. Range of g $\not\subset$ domain of g oe	B1													
5(a)	$\frac{10}{2}(6 + 9d) = 4 \times \frac{5}{2}(6 + 4d)$	M2	M1 for either sum correct												
	$d = 6$	A1													

Question	Answer	Marks	Guidance
5(b)	$ar^2 = a + 4(ar - a)$	M1	For $ar = a + d$, $ar^2 = a + 4d$ or for $ar^2 = a + k(ar - a)$ ($k > 2$) with a correct attempt to eliminate d
	$r^2 - 4r + 3 = 0$	A1	
	$(r - 3)(r - 1) = 0$ or formula	M1	Dep for solving <i>their</i> 3-term quadratic
	$r = 3$ only	A1	
	Alternative method		
	$\frac{a+d}{a} = \frac{a+4d}{a+d}$	(M1)	
	$\Rightarrow d = 2a$	(A1)	
	$r = \frac{a+2a}{a}$	(M1)	Dep
6	$r = 3$	(A1)	
	$x^2 - 5x = 6 \Rightarrow x^2 - 5x - 6 = 0$		
	$(x - 6)(x + 1) = 0$	M1	or $x = \frac{5 \pm \sqrt{25 - 4 \times (-6)}}{2}$
	$x = 6$ or -1	A1	
	$x^2 - 5x = -6 \Rightarrow x^2 - 5x + 6 = 0$	M1	
	$(x - 3)(x - 2) = 0$	M1	or $x = \frac{5 \pm \sqrt{25 - 4 \times 6}}{2}$
	$x = 3$ or 2	A1	
	Alternative method		
	$(x^2 - 5x)^2 = 36$	(M1)	
	$x^4 - 10x^3 + 25x^2 - 36 = 0$	(A1)	
	$(x - 3)(x - 2)(x - 6)(x + 1) = 0$	(M1)	Attempt to factorise into 4 brackets
	$x = -1$ or 2 or 3 or 6	(A1A1)	A1 for any two

Question	Answer	Marks	Guidance
7(a)	$\frac{e^{1-3x}(\cos x - \sin x) - (\sin x + \cos x) \times (-3)e^{1-3x}}{(e^{1-3x})^2}$	4	B1 for $-3e^{1-3x}$ B1 for $\cos x - \sin x$ M1 for differentiation of quotient or equivalent correct product A1 FT on <i>their</i> $-3e^{1-3x}$ and <i>their</i> $\cos x - \sin x$
7(b)	$\int (\sec^2 3x) dx$	M1	Use of $\sec^2 3x = 1 + \tan^2 3x$
	$\frac{1}{3} \tan 3x + c \text{ oe}$	2	A1 if $+ c$ omitted
8	$(1-6y)^2 - 8y^2 = -2y + 12y^2 \text{ oe}$	M1	Equation in one variable only, in any form. Allow one arithmetical error
	$16y^2 - 10y + 1 [= 0]$	A1	
	$(2y - 1)(8y - 1) [= 0]$	M1	Attempt to factorise or use formula for <i>their</i> 3-term quadratic
	$y = \frac{1}{2}, x = -2$ $y = \frac{1}{8}, x = \frac{1}{4}$	2	Correct pairs seen or implied A1 for any two correct values
9(a)	As $v-t$ graph but negative portion reflected in x -axis	B1	
9(b)	Straight line with positive gradient	B1	
	Straight line starting at O	B1	

Question	Answer	Marks	Guidance
10	Equation of normal $y = -x + 3$ soi	2	M1 for a correct attempt to obtain an equation of the normal using calculus and perpendicular gradient
	$-x + 3 = x^2 - 9x + 18$ or $x^2 - 9x + 18 = 0$	M1	
	$x = 3$ [or 5]	A1	
	$\int_3^5 [(-x+3) - (x^2 - 9x + 18)] dx$ $\pm \left[\left(\frac{-x^2}{2} + 3x \right) - \left(\frac{x^3}{3} - \frac{9x^2}{2} + 18x \right) \right]_3^5$ or $\pm \left[-\frac{x^3}{3} + 4x^2 - 15x \right]_3^5$	4	M1 for attempt to subtract and integrate Allow if order is reversed FT on <i>their</i> normal equation for M1 A1 for $\frac{x^3}{3} - \frac{9x^2}{2} + 18x$ soi A1 for $\frac{-x^2}{2} + 3x$ soi Ignore limits at this stage
	$\pm \left(-\frac{125}{3} + 100 - 75 - (-9 + 36 - 45) \right) \text{ oe}$	2	M1 dep for attempt to use <i>their</i> limits Allow one arithmetic error for M1 May include area of triangle rather than area using the normal
	$\frac{4}{3}$	A1	Curve – line gives $-\frac{4}{3}$, final answer must be $\frac{4}{3}$
Alternative method			
	$x^2 - 9x + 18 = 0$	(M1)	
	$x = 3$ [or 5]	(A1)	
	$\int_3^5 (x^2 - 9x + 18) dx$ $= \left[\frac{x^3}{3} - \frac{9x^2}{2} + 18x \right]_3^5$	(3)	M1 for attempt to integrate A1 for one term correct soi Ignore limits at this stage
	$\pm \left(\frac{125}{3} - \frac{225}{2} + 90 - \left(9 - \frac{81}{2} + 54 \right) \right) \text{ soi}$	(2)	M1 dep for attempt to use <i>their</i> limits Allow one arithmetic error for M1
	Area of triangle = $\frac{1}{2} \times 2 \times 2$	(2)	M1 for a complete method
	Shaded area = $\frac{10}{3} - 2 = \frac{4}{3}$	(2)	M1 for a difference between <i>their</i> areas Must be positive for A1

Question	Answer	Marks	Guidance
11(a)	$\frac{n!}{(n-4)!} = \frac{(2n)!}{6(2n-3)!}$	B1	
	$n(n-1)(n-2)(n-3) = \frac{2n(2n-1)(2n-2)}{6}$	2	Dep on previous B1 B1 for simplification of one side
	e.g. $3(n-2)(n-3) = 2(2n-1)$ $3n^2 - 19n + 20 = 0$ AG	B1	Dep on all previous B marks With sufficient detail shown
11(b)	$(3n-4)(n-5) = 0$ $n = 5$ only	2	M1 for attempt to solve the given equation
12	$\overrightarrow{AB} = \begin{pmatrix} 7 \\ 4 \end{pmatrix} - \begin{pmatrix} 1 \\ 7 \end{pmatrix} \quad \left[= \begin{pmatrix} 6 \\ -3 \end{pmatrix} \right]$	M1	
	$\overrightarrow{AC} = k \begin{pmatrix} 1 \\ 2 \end{pmatrix} - \begin{pmatrix} 1 \\ 7 \end{pmatrix} \quad \left[= \begin{pmatrix} k-1 \\ 2k-7 \end{pmatrix} \right]$	M1	
	Use of $\overrightarrow{AB} \parallel \overrightarrow{AC}$ or $\overrightarrow{AC} = m\overrightarrow{AB}$ to obtain either $\frac{k-1}{6} = \frac{2k-7}{-3}$ or $k-1 = 6m$ and $2k-7 = -3m$ oe	M1	Form equation without vectors FT their \overrightarrow{AB} and \overrightarrow{AC}
	$k = 3$	A1	
	$\overrightarrow{AC} = \begin{pmatrix} 3-1 \\ 2 \times 3 - 7 \end{pmatrix} = \begin{pmatrix} 2 \\ -1 \end{pmatrix}$ or $m = \frac{1}{3}$	M1	Substitute their k into their \overrightarrow{AC}
	$AC : AB = 1 : 3$ or $\frac{1}{3}$	A1	