



Cambridge International AS & A Level

CANDIDATE
NAME
CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



PHYSICS

9702/21

Paper 2 AS Level Structured Questions

October/November 2025

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 60.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Any blank pages are indicated.

Data

acceleration of free fall	$g = 9.81 \text{ m s}^{-2}$
speed of light in free space	$c = 3.00 \times 10^8 \text{ m s}^{-1}$
elementary charge	$e = 1.60 \times 10^{-19} \text{ C}$
unified atomic mass unit	$1 \text{ u} = 1.66 \times 10^{-27} \text{ kg}$
rest mass of proton	$m_p = 1.67 \times 10^{-27} \text{ kg}$
rest mass of electron	$m_e = 9.11 \times 10^{-31} \text{ kg}$
Avogadro constant	$N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$
molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Boltzmann constant	$k = 1.38 \times 10^{-23} \text{ J K}^{-1}$
gravitational constant	$G = 6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$
permittivity of free space	$\varepsilon_0 = 8.85 \times 10^{-12} \text{ F m}^{-1}$ $(\frac{1}{4\pi\varepsilon_0} = 8.99 \times 10^9 \text{ m F}^{-1})$
Planck constant	$h = 6.63 \times 10^{-34} \text{ J s}$
Stefan–Boltzmann constant	$\sigma = 5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$

Formulae

uniformly accelerated motion	$s = ut + \frac{1}{2}at^2$ $v^2 = u^2 + 2as$
hydrostatic pressure	$\Delta p = \rho g \Delta h$
upthrust	$F = \rho g V$
Doppler effect for sound waves	$f_o = \frac{f_s v}{v \pm v_s}$
electric current	$I = Anvq$
resistors in series	$R = R_1 + R_2 + \dots$
resistors in parallel	$\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2} + \dots$





BLANK PAGE

DO NOT WRITE IN THIS MARGIN



1 (a) Define acceleration.

.....
.....

(b) A rocket is launched vertically from the surface of the Earth.

Fig. 1.1 shows the variation of the velocity of the rocket with time for the first 20s after its launch.




Fig. 1.1

(i) Determine the acceleration of the rocket.

$$\text{acceleration} = \dots \text{ms}^{-2} \quad [1]$$

(ii) Show that the height of the rocket above the surface of the Earth at a time of 20s after launch is 3.2km.

[2]



(c) The mass of the rocket in (b) is $2.9 \times 10^6 \text{ kg}$. Assume that this mass remains constant.

For this rocket, from launch to its height at a time of 20 s after launch:

(i) calculate the gain in gravitational potential energy ΔE_P

$$\Delta E_P = \dots \text{ J} [2]$$

(ii) calculate the gain in kinetic energy ΔE_K

$$\Delta E_K = \dots \text{ J} [2]$$

(iii) determine the average power output of the rocket engines. Assume that resistive forces are negligible.

$$\text{power} = \dots \text{ W} [2]$$

[Total: 10]



2 (a) (i) Define pressure.

.....
.....

[1]

(ii) Explain how hydrostatic pressure results in an upthrust force acting on a solid object immersed in a liquid.

.....
.....
.....
.....

[2]

(b) A small steel ball of radius r and mass m falls vertically at terminal speed v through oil.

The viscous drag force D that acts on the ball is given by

$$D = 6\pi\eta rv$$

where η is a property of the oil called its viscosity.

(i) On Fig. 2.1, draw labelled arrows from the ball to show the directions of the **three** forces that act on the ball as it falls.



Fig. 2.1

[3]





(ii) Determine the SI base units of η .

base units [2]

(c) The oil in (b) has a density of 920 kg m^{-3} and a viscosity of 4.7 in SI units.

The steel ball has a mass of $2.4 \times 10^{-3} \text{ kg}$ and a radius of $4.2 \times 10^{-3} \text{ m}$.

(i) Show that the upthrust force acting on the ball is $2.8 \times 10^{-3} \text{ N}$.

[1]

(ii) Determine the terminal speed v of the ball.

$v =$ ms^{-1} [3]

[Total: 12]



3 A wire has length L and cross-sectional area A . The wire is made from a metal that has Young modulus E and resistivity ρ .

(a) Define the Young modulus of a material.

.....
.....

[1]

(b) (i) State an expression, in terms of some or all of L , A , E and ρ , for the resistance R_0 of the wire.

$$R_0 = \dots \quad [1]$$

(ii) Show that the spring constant k_0 of the wire is given by

$$k_0 = \frac{EA}{L}.$$

[2]

(c) The wire is stretched, within the limit of proportionality, by a tensile force F . Assume that any changes in the cross-sectional area of the wire are negligible.

(i) On Fig. 3.1, sketch the variation with F of the resistance R of the wire.



Fig. 3.1

[1]



(ii) On Fig. 3.2, sketch the variation with F of the spring constant k of the wire.



Fig. 3.2

[1]

(d) Copper has a resistivity of $1.8 \times 10^{-8} \Omega \text{ m}$ and a Young modulus of $1.3 \times 10^{11} \text{ Pa}$.

A copper wire of diameter 1.6 mm has a resistance of 0.034Ω .

(i) Show that the length of the wire is 3.8 m.

[1]

(ii) Use the equation in (b)(ii) to determine the spring constant of the wire.

spring constant = Nm^{-1} [2]

[Total: 9]



4 (a) State what is meant by diffraction of a wave.

.....
.....
.....

[2]

(b) A beam of vertically polarised light of wavelength 540 nm is incident normally on a diffraction grating, as shown in Fig. 4.1.




Fig. 4.1 (not to scale)

The diffraction grating has a line spacing of 5.0×10^{-6} m.

The light transmitted by the diffraction grating illuminates a circular screen. The diffraction grating is at the centre X of the circle.

The central bright fringe is formed at point O on the screen and has intensity I_0 .

P is a point on the screen where the line XP is at a variable angle θ to the line XO. The intensity I of light on the screen at P varies with θ .

(i) Show that the angle θ at which the first-order bright fringe is formed is 6.2° .

[2]



(ii) Determine the value of θ at which the second-order bright fringe is formed.

$$\theta = \dots \text{ } ^\circ [1]$$

(iii) On Fig. 4.2, sketch the variation of the intensity I with θ for values of θ from -15° to $+15^\circ$.




Fig. 4.2

[3]

(c) A polarising filter is placed in the path of the light beam that is incident on the diffraction grating in Fig. 4.1. The transmission axis of the filter is at 45° to the vertical.

Suggest how the variation of intensity with θ for the light on the screen compares with the answer in (b)(iii).

.....
.....
.....

[2]

[Total: 10]



5 (a) State Kirchhoff's first law.

.....
.....

[1]

(b) Fig. 5.1 shows a circuit containing a thermistor T that has a negative temperature coefficient.




Fig. 5.1

(i) The thermistor has resistance R_0 at a temperature of 0°C .

On Fig. 5.2, sketch a possible variation of the resistance of the thermistor with temperature between 0°C and 100°C .




Fig. 5.2

[2]



(ii) With reference to the current in the cell, explain why the current in **resistor R** decreases with increasing temperature of the thermistor.

.....
.....
.....
.....
.....

[3]

(c) The electromotive force (e.m.f.) E of the cell in Fig. 5.1 is 1.50V. The internal resistance r of the cell is 0.12Ω .

Resistor R has a resistance of 6.00Ω .

At a particular temperature of the thermistor, the current in R is 0.200A.

For this temperature of the thermistor, determine:

(i) the current in the cell

current =A [2]

(ii) the resistance of the thermistor.

resistance = Ω [2]

[Total: 10]



6 The nuclide ${}^3_1\text{H}$ is an isotope of hydrogen that is called tritium.

(a) (i) Determine the numbers of protons, neutrons and electrons in a neutral atom of tritium.

number of protons =

number of neutrons =

number of electrons =


[2]

(ii) Draw a labelled diagram to represent a simple model of the arrangement of the protons, neutrons and electrons in a tritium atom.

[2]

(b) Tritium is radioactive and undergoes β^- decay to form an isotope of helium (He). Gamma radiation is not emitted during this decay.

(i) Complete the equation to represent the radioactive decay of tritium.



[2]

(ii) State the name of particle X.

..... [1]



(c) Determine the quark composition of a tritium nucleus.

..... [2]

[Total: 9]





BLANK PAGE

DO NOT WRITE IN THIS MARGIN

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

